

CERN-TH/97-354

KUL-TF-97/35

hep-th/9712135

D0-Branes as Instantons in $D = 4$ Super Yang-Mills Theories ¹

J. L. F. Barbón [†] and A. Pasquinucci [‡]

[†] *Theory Division, CERN*
CH-1211 Geneva 23, Switzerland
`barbon@mail.cern.ch`

[‡] *Instituut voor theoretische fysica, K.U. Leuven*
Celestijnenlaan 200D, B-3001 Leuven, Belgium

Abstract: We discuss the role of D0-branes as instantons in the construction of $SU(N)$ Super Yang-Mills and Super QCD theories in four space-time dimensions with D4-, D6- and NS-branes.

CERN-TH/97-354

December, 1997

¹ To appear in the proceedings of the conference “*Quantum Aspects of Gauge Theories, Supersymmetry and Unification*”, held at Neuchâtel University, Neuchâtel, Switzerland, 18–23 September 1997.

D0-Branes as Instantons in $D = 4$ Super Yang-Mills Theories

J. L. F. Barbón¹ and A. Pasquinucci²

¹ Theory Division, CERN

CH-1211 Geneva 23, Switzerland

² Instituut voor theoretische fysica, K.U. Leuven

Celestijnenlaan 200D, B-3001 Leuven, Belgium

Abstract: We discuss the role of D0-branes as instantons in the construction of $SU(N)$ Super Yang-Mills and Super QCD theories in four space-time dimensions with D4-, D6- and NS-branes.

The basic construction of SYM instantons on a Dp -brane world-volume, is as bound states with $D(p-4)$ -branes [1]. In the case of the construction of refs. [2] (see also [3]), the $SU(n_c)$ SYM theory is described by n_c D4-branes, so we must consider D0-branes bound to D4-branes. The five-dimensional world-volume of the D4-branes is then compactified to the four-dimensional physical space and, to make a bound D0-brane describe an instanton in the four-dimensional physical space, the D0-brane Euclidean-time world-line must lie along the compactified direction of the D4-brane. Although the details of such a reduction depend on the amount of supersymmetry ($N = 4, 2, 1, 0$) that we want to preserve in space-time, the basic properties of the instanton moduli space only depend on the geometry of D0-D4-branes bound states, which we now review.

The collective dynamics of the instantons is specified by a point action and an appropriate measure over instanton moduli space. In the present D0-branes construction, both ingredients appear as the dimensional reduction down to zero dimensions of the D0-brane one-dimensional collective dynamics. In particular, the instanton moduli space as a hyper-Kähler quotient appears as a Higgs branch of the world-line theory of the D0-branes. The basic terms in the world-line action are the static Born-Infeld action of the D0-brane

$$S_{BI} = \tau_0 \int \sqrt{\det h_{\text{induced}}} \longrightarrow M_{D0} \int d\tau, \quad (1)$$

and a source term coming from the Chern-Simons couplings on the D4-branes world-volume

$$S_{\text{source}} = i \frac{\mu_4}{2} \int_{\Sigma_{4+1}} A_{D0}^{RR} (2\pi\alpha')^2 F \wedge F, \quad (2)$$

which relates the RR photon of the type-IIA string theory to the theta parameter of the four-dimensional instanton. The basic D0-brane action (1) leads to a fit of the effective length of the world-line, in order to reproduce the instanton action $S_{\text{inst}} = 8\pi^2/g^2$. Indeed from the Born-Infeld action of the D4-brane gauge theory, upon dimensional reduction in the world-line direction (say, the x^6 direction), on an interval of length L_6 , we have

$$\tau_4 \int_0^{L_6} d\tau \text{Tr} \sqrt{\det(1 + 2\pi\alpha' F)} \rightarrow -\frac{\tau_4}{4} L_6 (2\pi\alpha')^2 \text{Tr} F^2 \equiv -\frac{1}{4g_{YM}^2} \text{Tr} F^2. \quad (3)$$

Using $\tau_p^{-1} = g_s \sqrt{\alpha'} (2\pi \sqrt{\alpha'})^p$ and $M_{D0} = \tau_0$ we find agreement since $8\pi^2/g^2 \equiv M_{D0} L_6 = 8\pi^2/g_{YM}^2$. Also, for a constant one-form A_{D0}^{RR} , the five-dimensional integral (2) is localized on the D0-brane world-line, and we have $S_{\text{source}} = -i \frac{\theta}{T_6} \int d\tau$.

The zero-mode structure of the instantons comes from the non-trivial world-volume dynamics of the D0-branes, as given by the massless limit of the corresponding open strings. There are non-trivial gauge interactions specifying the dynamics of 0-0 strings and 0-4 strings, in the general case where we have k D0-branes bound to n_c D4-branes. The 0-0 sector is just the dimensional reduction down to one dimension of ten-dimensional SYM with $U(k)$ gauge group, corresponding to a system of k D0-branes, and has $N = 4$ supersymmetry in four-dimensional notation. The 0-4 sector breaks half of these unbroken 16 supercharges, leading to an action for the 0-4 strings with $N = 2$ supersymmetric couplings.

To be more explicit, let us parametrize the different degrees of freedom in four-dimensional superfield notation, i.e. we define the world-line action by dimensional reduction from an $N = 2$ model in four dimensions, which can be obtained as a T-dual configuration of D3-branes inside D7-branes of type IIB string theory. This auxiliary four-dimensional space is along the (x^6, x^7, x^8, x^9) directions. We denote by M the physical Minkowski space-time (x^0, x^1, x^2, x^3) and after a Wick rotation, we shall work with complex coordinates $z = x^0 + ix^1$, $\tilde{z} = x^2 + ix^3$ and $w = x^4 + ix^5$. The 0-0 matrix coordinates, representing generalized translational degrees of freedom of the k D0-branes, are described by a set of chiral superfields $\Phi_{00} \equiv (W, Z, \tilde{Z})$, representing displacements in the corresponding coordinates, and by the dimensionally reduced vector superfield $V_{00} \sim (A_6, A_7, A_8, A_9)$, plus superpartners, all in the adjoint representation of $U(k)$. We will not consider excitations or deformations of the field A_7 , and we will gauge the longitudinal A_6 to zero (recall that the D0-branes world-lines lie in the x^6 direction). Finally, the 0-4 and 4-0 sectors give rise to hypermultiplets, represented by chiral superfields $\Phi_{04} = H, \Phi_{40} = \tilde{H}$, transforming in the $(n_k, \bar{n}_c), (\bar{n}_k, n_c)$ of $U(k) \times U(n_c)$ respectively. The complete superpotential contains an $N = 4$ interaction between 0-0 fields:

$$\mathcal{W}_{0-0} = \frac{\sqrt{2}}{g_s} \frac{1}{3!} \epsilon_{ijk} \text{Tr } \Phi_i [\Phi_j, \Phi_k] \quad (4)$$

and an $N = 2$ preserving superpotential for the “quarks”:

$$\mathcal{W}_{0-4} = \frac{\sqrt{2}}{g_s} (\tilde{H} W H - H W' \tilde{H}), \quad (5)$$

where we have introduced a mass term inherited from the superpotential in the D4-branes, i.e. the 4-4 strings, coupling the $SU(n_c)$ adjoint $W'_{44} = X'_4 + iX'_5$ giving the position of D4-branes. The $N = 2$ supersymmetry on the D0-brane world-line implies that $[W', W'^{\dagger}] = 0$, i.e. the W' field lies along the flat directions of the D4-brane theory, and can be taken as diagonal matrices by an $SU(n_c)$ rotation¹ $W'^b_a = w'_a \delta^b_a$. This lagrangian also appears in various M-atrity theory applications [5].

The instanton moduli space is then obtained as a maximal Higgs phase of this theory on the D0-brane world-line, where we consider the most general vacuum expectation values of the hypermultiplets $H, \tilde{H}$ and the adjoint fields $Z, \tilde{Z}$. In this paper we will be mainly concerned with properties of the single instanton sector, or equivalently, of the dilute gas limit of instantons. In this case the overlap between the instantons can be neglected and the moduli space admits a much simpler description. Indeed, well separated D0-branes are characterized by the complete diagonalization of all the adjoint 0-0 fields corresponding to the space-time positions, i.e. we consider $Z, \tilde{Z}$ diagonal generic matrices, which spontaneously break $U(k)$ to the Abelian subgroup $U(1)^k$. It is intuitively clear that this produces just k copies of the single instanton moduli space. In this subspace, the $4k$ position degrees of freedom $z_i, \tilde{z}_i$ are completely decoupled from the rest of the variables.

The interesting branch of solutions is the one with maximal Higgsing of the hypermultiplets, with $\text{diag}(W) = \text{diag}(W') = 0$ (all D0 and D4-branes on top of each other in the (x^4, x^5) plane). Subtracting the $2k$ F -equations and the k D -equations from the total $4kn_c$ real components in the scalars $\phi_h, \tilde{\phi}_h$,

¹The brane construction of the full gauge theory produces naively a $U(n_c)$ group. Only the $SU(n_c)$ subgroup is relevant to the instanton discussion, of course. In fact, on the Coulomb branch of the D4-branes world-volume, the extra $U(1)$ is frozen, since fluctuations of the trace of W' have infinite norm [4].

and further dividing by the $U(1)^k$ gauge symmetry, we find a total of $4kn_c - 2k - k - k = 4k(n_c - 1)$ real parameters. Adding the $4k$ degrees of freedom coming from the eigenvalues of Z and $\tilde{Z}$, representing the translation moduli in the physical dimensions, we end up with the correct $4kn_c$ dimensional bosonic moduli space of YM instantons.

The basic idea in this microscopic model for YM instantons is then the Kaluza–Klein truncation of the D4 gauge theory with 16 supercharges, down to a four-dimensional theory, together with a zero-mode truncation of the world-line dynamics of the D0-branes.

In the simplest case, upon reduction on a circle of length L_6 , we get $N = 4$ SYM in four dimensions. On the relevant moduli space of the D0-branes, we are instructed to keep only the zero modes. For k D0-branes, the world-line has a set of $d = 2kn_c$ free superfields (ξ_s, η_s) , in four-dimensional notation, i.e. we have an action

$$S_{D0} = kM_{D0} \int_0^{L_6} d\tau - ik\theta + \frac{1}{g_s} \int_0^{L_6} d\tau \sum_{s=1}^d \left(|\dot{\xi}_s|^2 + i\bar{\eta}_s \dot{\eta}_s \right) + \text{massive} . \quad (6)$$

Each Weyl fermion η_s has two complex or four real components off-shell so that, when the one-instanton contribution to the path-integral is dominated by the classical static trajectory $\dot{\xi}_s = \dot{\eta}_s = 0$, we get a factor

$$Z_{\text{inst}} = \int \prod_s d\bar{\eta}_s d\eta_s d\mu(\xi_s) e^{-kM_{D0}L_6 + ik\theta} = 0^{4d} \text{Vol}(\mathcal{M}) e^{-\frac{8\pi^2 k}{g^2} + ik\theta} \quad (7)$$

where $M_{D0}L_6 \equiv 8\pi^2/g^2$. In this expression, the bosonic measure $d\mu(\xi)$ for collective coordinates contains zero-mode Jacobians which are most easily derived in the field theoretical framework. The corresponding integral gives the formally divergent volume of moduli space, $\text{Vol}(\mathcal{M})$. The symbol 0^{4d} denotes the number of fermionic zero modes of the instanton, to be saturated by fermion sources. We see clearly that the number of zero modes corresponds to the total number of independent anticommuting collective coordinates, which in turn coincide with the total number of off-shell fermion components on the D0-brane world-line. The corresponding number of fermion zero-modes is $4d = 8kn_c$, which is indeed the right one for $SU(n_c)$ SYM in the k -instanton sector. In field theory, each gluino system gives $2kn_c$ zero modes, and we have four independent gluino systems in $N = 4$ SYM. In the D0-D4 branes construction, the $8k$ zero modes associated to the $Z, \tilde{Z}$ superfields are interpreted in the four-dimensional SYM theory as supersymmetry zero modes. On the other hand, $8k$ out of the $8k(n_c - 1)$ zero modes associated to the hypermultiplets $H, \tilde{H}$ are interpreted in space time as superconformal zero modes, while the remaining $8k(n_c - 2)$ zero modes do not have an obvious symmetry interpretation, since they arise as 't Hooft zero modes associated to doublets with respect to the $SU(2)$ subgroup of the gauge group where the instanton sits.

According to the constructions of refs. [2], models with $N = 2$ and $N = 1$ supersymmetry can be obtained upon compactification on a segment of length L_6 , bounded by two NS5-branes. One of them, denoted NS, extends in (M, w) , and the other, NS', possibly rotated, extends in (M, Π) , with Π a plane in the (w, v) space. In the case of $N = 2$ supersymmetry the NS and NS' branes are parallel, whereas for $N = 1$ the plane Π is obtained from the w -plane by an $SU(2)$ rotation. For the purposes of the D0-branes dynamics, the role of the NS5-branes is just to project out some fermionic zero modes, namely, those zero modes among the total of $8kn_c$ anticommuting coordinates $\eta_s, \bar{\eta}_s$ that overlap with broken supersymmetries of the brane configuration in M . These zero modes correspond really to Goldstone fermions and are not normalizable in the supersymmetric vacuum of M . Further, we assume that no projection is imposed by the NS5-branes on the bosonic collective coordinates of the D0-branes. Indeed to the extent that they are bound to the D4-branes, they can freely move in (the Euclidean version of) M , and furthermore the Higgs branch collective coordinates do not feel any geometric constraint as a result of the presence of the NS5-branes.

In order to see how the fermion projections work, we recall that the unbroken supersymmetry on a brane world-volume is best represented in M-theory language in terms of the projectors $P_p = \frac{1}{2}(1 + i\Gamma(\Sigma_{p+1}))$, where $\Gamma(\Sigma_{p+1})$ is the product of Dirac matrices in the world-volume directions, with the

understanding that p -branes whose M-theory description involves a wrapped or boosted M-brane in the eleventh direction include a factor of Γ_{11} . So, in terms of a 32-dimensional spinor Ψ_{32} , the 16 Goldstone fermions carried by a free D0-brane are given by $(1 - P_{D0})\Psi_{32}$. Now, according to the above prescription, we should keep only those zero modes that have no overlap with the Goldstone fermions on the NS-brane world-volume. The remaining zero modes from the 0-0 sector of a general free D0-brane are then $\eta_{z,\bar{z},v,w} = (1 - P_{D0})P_{NS}P_{NS'}\Psi_{32}$. Using the explicit form of the projectors, one easily sees that each NS-brane divides by half the number of zero modes, so that we have 8 zero modes in the $N = 2$ configuration and 4 zero modes in the $N = 1$ configuration.

The previous considerations apply to the situation of a free D0-brane. In the case of a D0-D4-branes bound state we have two modifications. First, as stated before, on the Higgs branch of bound states the fermions ψ_v, ψ_w are lifted, and the number of zero modes coming from the 0-0 sector is again divided by half. Second, we have a new set of degrees of freedom, namely the $\psi_h, \bar{\psi}_h$ from the 0-4 and 4-0 sectors. Those are constructed from the zero modes of the world-sheet fermions ψ_0^μ for $\mu = 4, 5, 6, 7, 8, 9$ (i.e. in the NN + DD directions of the D0-D4-brane system), in the Ramond sector. Indeed, there are $\frac{1}{2}2^{6/2} = 4$ field components (off shell), after GSO projection, in the 0-4 sector, that is a Weyl fermion. In terms of the complex combinations $\Gamma_u = \frac{1}{2}(\Gamma_6 + i\Gamma_7)$, $\Gamma_w = \frac{1}{2}(\Gamma_4 + i\Gamma_5)$, $\Gamma_v = \frac{1}{2}(\Gamma_8 + i\Gamma_9)$, the chirality operator reads

$$\Gamma_{11} = i\Gamma_0\Gamma_1 \cdots \Gamma_9 = -(1 - 2N_z)(1 - 2N_{\bar{z}})(1 - 2N_u)(1 - 2N_v)(1 - 2N_w), \quad (8)$$

with $N_v \equiv \Gamma_v^\dagger \Gamma_v$, etc., the fermion occupation numbers taking values 0 or 1. Acting on the $2^{6/2} + 2^{6/2}$ states in 0-4 and 4-0 sectors, built as polynomials $P(\Gamma_u^\dagger, \Gamma_v^\dagger, \Gamma_w^\dagger)|0\rangle$, the GSO projection reduces to $-(1 - 2N_u)(1 - 2N_v)(1 - 2N_w) = +1$. On the same set, the projection imposed by the NS brane is $+1 = i\Gamma_{NS} = i\Gamma_0\Gamma_1 \cdots \Gamma_4\Gamma_5 = -(1 - 2N_w)$. So, on this subspace the NS projector is $P_{NS} = N_w$, and the analogous projector for the rotated NS'-brane is $P_{NS'} = N_v$.

We see that, in all cases, the effect of each NS-brane is to divide by 2 all the bulk zero modes democratically. This amounts to a total factor of $1/4$ when the NS' brane is a rotation of the NS-brane into the (x^8, x^9) plane. So, if N denotes the amount of space-time supersymmetry of the brane configuration, the resulting instanton leaves $2N$ supercharges unbroken, out of the total $4N$ that are linearly realized in the vacuum of the space-time theory. The total number of fermionic zero-modes is given by:

$$\mathcal{N}_{f.z.m.} = \frac{N}{4} \cdot 4d = \frac{N}{4} \cdot 8k + \frac{N}{4} \cdot 8k(n_c - 1) = 2Nkn_c, \quad (9)$$

the correct number, where we have already subtracted the massive degrees of freedom in the mixed Higgs-Coulomb branch, which defines the instanton moduli space, and we have split the zero modes between those corresponding to the $2N$ broken supercharges, and those of a fixed instanton, coming from the hypermultiplet degrees of freedom.

Introducing flavour brings in new subtleties. Now we have n_f D6-branes extended in (M, v, x^7) , and localized at definite values of x^6 . Since the D0-branes world-line extends in x^6 , the local D0-D6 intersections are instantonic! This means that the 0-6 and 6-0 strings must be quantized with boundary conditions such that $\nu = d_{ND} + d_{DN} = 8$ (the sum of ND and DN directions). Thus the 0-6 sector is T-dual to a D1-D9 sector, and massless fermionic degrees of freedom appear localized at the D0-D6 intersections. Carrying out the analysis [6], one finds the required $2kn_f$ fermionic zero modes associated to the quarks in the space-time description. Since these zero modes are localized at intermediate points in the D0 world-line, they are insensitive to the projections imposed by the NS5-branes at the end-points. Therefore, the number of flavour zero modes depends only on n_f , but not on the amount of space-time supersymmetry, in beautiful correspondence with the field theory result.

Up to now we have considered the case of exact instantons in super Yang-Mills theory and showed that it can be described by a system of k D0-branes bound to n_c D4-branes. In particular this means that the D0-branes and the D4-branes are all located at the same point in the $(x^4, x^5, x^7, x^8, x^9)$ plane. The geometrical operation of moving the D4-branes apart corresponds, as intuition would tell, to describing constrained instantons [7]. The relevant coupling in the D0-brane world-line theory, which is sensitive to

the D4-branes motion is (see eq. (5))

$$\mathcal{L}_{04-44-40} = \frac{\sqrt{2}}{g_s} \int d^2\theta \Phi_{04} \Phi_{44} \Phi_{40} + \text{h.c.} \quad (10)$$

This coupling is simply a mass term (the mass matrix given by $\phi_{44} \sim W'$ that describes the position of the D4-branes in the (x^4, x^5) plane) for the instanton moduli (in the notation of (5) $\Phi_{04} \sim \tilde{H}$ and $\Phi_{40} \sim H$). When considered in the Coulomb branch of the $N = 2$ configurations, it yields the familiar action of a constrained instanton:

$$S_{\text{constr}} = \frac{1}{g^2} (8\pi^2 + \rho^2 v^2) , \quad (11)$$

with $v \sim \langle \phi_{44} \rangle$ and $\rho^2 \sim \phi_{04} \phi_{40}$. At the same time, it lifts the fermion zero modes through the pairings $v \psi_{04} \psi_{40}$. For the fermions, the lifting of zero modes occurs homogeneously in the bulk of the D0-brane world-line, and therefore it is obviously compatible with the projections imposed by the NS-branes at the boundaries. In general, a homogeneous lifting pattern of this sort, appropriate to describe the Coulomb phase of $N = 2$ models, gives a mass of order v to d_c complex bosonic collective coordinates, and similarly, it lifts $4d_c$ real fermionic coordinates in the bulk, i.e. it induces a term in the bulk action of the form $\delta\mathcal{L} \sim v^2 \sum_{s=1}^{d_c} |\xi_s|^2 + v \sum_{s=1}^{d_c} \bar{\eta}_s \eta_s$, where we have rescaled the bare Yang-Mills coupling $g^2 \sim g_s/L_6$ into the collective coordinates ξ_s and η_s . Taking into account the projection at the boundaries of the world-line due to the NS- and NS'-branes, we have a partition function of the form

$$Z_{\text{inst}} \sim 0^{N(d-d_c)} \text{Vol}_c(\mathcal{M}) v^{\frac{Nd_c}{2}} e^{-\frac{8\pi^2}{g^2} + i\theta} \quad (12)$$

where $\text{Vol}_c(\mathcal{M})$ represents the bosonic “volume” of the moduli space, regulated with the Gaussian term $\exp(-v^2|\xi|^2)$, and the term $v^{Nd_c/2}$ comes from the fermionic integrals.² The complete power of v in the full instanton measure depends on the result of the bosonic integrals, of course. In general, we must view the couplings like eq. (10) as small perturbations over the instanton moduli space described in the introduction. This amounts to the requirement that the dimensionless effective expansion parameter ρv be small.

The important remark to make is that all zero modes due to the conformal symmetry and embedding angles, namely, all zero modes encoded in Φ_{04}, Φ_{40} , can be lifted in this way on the Coulomb branch of the $N = 2$ theory, by switching on Φ_{44} (i.e. splitting the D4-branes in the (x^4, x^5) plane). This amounts to generically lifting $2d_c = 4(n_c - 1)$ bosonic and $4(n_c - 1)$ fermionic zero modes. The remaining zero modes, 4 bosonic and 4 fermionic, correspond to the superfields from the 0-0 sector Z and $\tilde{Z}$, which remain uncoupled and therefore are not lifted. Since these collective coordinates are associated to the translations in the physical four dimensions, these fermion zero modes are naturally interpreted as due to the broken supersymmetries of the instanton, understood as a classical solution in the physical four-dimensional theory on the four-branes.

Using these ideas, one can postulate simple couplings at the intersections between D0-, D4- and D6-branes, which explain the pattern of zero-mode lifting appropriate to $N = 2$ and $N = 1$ Higgs branches. Analogous, even if not so detailed, considerations can be done also in the case of multi-instanton configurations beyond the dilute-gas approximations. We refer the reader to [6] for more details.

Although our discussion applies strictly to the regime of weak type-IIA coupling, it is interesting to ask ourselves what happens of the D0-branes we have been discussing in this paper, in the lifting to M-theory [4]. A free D0-brane of the type-IIA theory becomes a graviton Kaluza-Klein state carrying momentum along the circle in the eleventh direction, and electrically charged with respect to the $U(1)$ gauge field corresponding to the rotations around this circle. As discussed in refs. [8], the physics of such free D0-branes seems to have nothing to do with the field theory SYM models we are interested in. Whereas in the M-theory picture it could be less simple to decouple these modes, in the type-IIA construction we can

² Recall that N is the number of four-dimensional supersymmetries in the space-time configuration, and $d = 2n_c$ is the complex dimension of the bosonic instanton moduli space.

just declare that we do not consider free D0-branes. In this way, we are making an explicit distinction between the free D0-branes, and the D0-D4 branes bound states. Indeed this distinction is natural from the point of view of the type-IIA configuration, since there is no non-contractible circle where we could wrap a free D0-brane world-line to yield a semiclassical instanton contribution. On the other hand, the world-line of the bound D0-brane is just the Euclidean world-line of the D4-brane with which it is solidary. In other words, once the D0-branes are bound to the D4-branes, they must be considered as part of the dynamical data on the D4-branes world-volume.

Thus, the only D0-branes of interest are those that are bound to at least two D4-branes and are not allowed to become free. In the internal space, the D0-branes just completely adhere to the D4-branes since they differ only in the space-time directions where the D0-branes are at a point. In lifting to M-theory, only the internal space gets modified and from this point of view the bound D0-branes just completely merge in the D4-branes and they become part of the resulting five-brane. Actually one could see this from the opposite point of view: in extending the D4-branes to become an M-theory five-brane wrapped around the eleventh circle, one is effectively dressing them up with bound D0-branes. Further discussion of theta angle dependence in MQCD can be found in ref. [9].

Acknowledgements

This work is partially supported by the European Commission TMR programme ERBFMRX-CT96-0045 in which A.P. is associated to the Institute for Theoretical Physics, K.U. Leuven. A.P. would like to thank CERN for its hospitality while part of this work was carried out.

References

- [1] E. Witten, *Nucl. Phys.* **B460** (1996) 335, [hep-th/9510135](#); *Nucl. Phys.* **B460** (1996) 541, [hep-th/9511030](#). M.R. Douglas, “*Branes within Branes*”, [hep-th/9512077](#); “*Gauge Fields and D-branes*”, [hep-th/9604198](#). C. Vafa, *Nucl. Phys.* **B463** (1996) 435, [hep-th/9512078](#). M.R. Douglas and G. Moore, “*D-branes, Quivers and ALE Instantons*”, [hep-th/9603167](#).
- [2] A. Hanany and E. Witten, *Nucl. Phys.* **B492** (1997) 152, [hep-th/9611230](#). S. Elitzur, A. Giveon and D. Kutasov, *Phys. Lett.* **B400** (1997) 269, [hep-th/9702014](#).
- [3] J. de Boer, K. Hori, H. Ooguri, Y. Oz and Z. Yin, *Nucl. Phys.* **B493** (1997) 148, [hep-th/9702154](#). J.L.F. Barbón, *Phys. Lett.* **B402** (1997) 59, [hep-th/9703051](#). N. Evans, C.V. Johnson and A.D. Shapere, *Nucl. Phys.* **B505** (1997) 251, [hep-th/9703210](#). J.H. Brodie and A. Hanany, *Nucl. Phys.* **B506** (1997) 157, [hep-th/9704043](#). S. Elitzur, A. Giveon, D. Kutasov, E. Rabinovici and A. Schwimmer, *Nucl. Phys.* **B505** (1997) 202, [hep-th/9704104](#).
- [4] E. Witten, *Nucl. Phys.* **B500** (1997) 3, [hep-th/9703166](#).
- [5] M. Berkooz and M.R. Douglas, *Phys. Lett.* **B395** (1997) 196, [hep-th/9610236](#). O. Aharony, M. Berkooz, S. Kachru, N. Seiberg and E. Silverstein, “*Matrix Description of Interacting Theories in Six Dimensions*”, [hep-th/9707079](#).
- [6] J.L.F. Barbón and A. Pasquinucci, “*D0-Branes, Constrained Instantons and D=4 Super Yang-Mills Theories*”, [hep-th/9708041](#).
- [7] G. 't Hooft, *Phys. Rev.* **D14** (1976) 3432. Y. Frishman and S. Yankielowicz, *Phys. Rev.* **D19** (1979) 540. I. Affleck, *Nucl. Phys.* **B191** (1981) 429.
- [8] E. Witten, “*Branes And The Dynamics Of QCD*”, [hep-th/9706109](#). A. Hanany, M.J. Strassler and A. Zaffaroni, “*Confinement and Strings in MQCD*”, [hep-th/9707244](#).
- [9] J.L.F. Barbón and A. Pasquinucci, “*Softly Broken MQCD and the Theta Angle*”, [hep-th/9711030](#).